

Protocol for the embedding of mammalian tissues (murine skin, murine liver) using the LEICA EM Tissue Processor

I. Fischer and S. Reipert, Institute of Biochem. & Mol. Cell Biol., Vienna Biocenter, Dr. Bohr-Gasse 9, A-1030 Vienna

- Perfusion with 2.5% Paraformaldehyde + 0.5% Glutaraldehyde in PBS.
- Cutting the tissue in small pieces (1 mm³).
- Fixation in 3% Glutaraldehyde in Sorensen´s over night at 4°C (tissue already placed in TP baskets).

	Reagent	Time	Temp.	Agitation
1	1.5 % OsO ₄	90 min	R.T.	3
2	Sorensen´s	5 min	R.T.	3
3	Sorensen´s	5 min	R.T.	3
4	Sorensen´s	5 min	R.T.	3
5	Sorensen´s	5 min	R.T.	3
6	Ethanol 30 %	5 min	R.T.	3
7	Ethanol 30 %	10 min	R.T.	3
8	Ethanol 50 %	5 min	R.T.	3
9	Ethanol 50 %	10 min	R.T.	3
10	Ethanol 70 %	5 min	R.T.	3
11	Ethanol 70 %	10 min	R.T.	3
12	Ethanol 95 %	5 min	R.T.	3
13	Ethanol 95 %	10 min	R.T.	3
14	Ethanol 100 %	5 min	R.T.	3
15	Ethanol 100 %	5 min	R.T.	3
16	Ethanol 100 %	5 min	R.T.	3
17	Ethanol 100 %	10 min	R.T.	3
18	Propylene Oxide	5 min	R.T.	3
19	Propylene Oxide	5 min	R.T.	3
20	Propylene Oxide	10 min	R.T.	3
21	Agar 100: Propylene Oxide 1:2	15 min	R.T.	3
22	Agar 100: Propylene Oxide 1:1	30 min	R.T.	3
23	Agar 100: Propylene Oxide 2:1	45 min	R.T.	3

- Infiltration in pure epoxy resin (Agar100) over night.
- Fresh resin and polymerisation for 24 h at 60 °C.